

# K<sub>2</sub><sup>\*</sup>(1980)

$$I(J^P) = \frac{1}{2}(2^+)$$


Needs confirmation.

## K<sub>2</sub><sup>\*</sup>(1980) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>1995<sup>+60</sup><sub>-50</sub> OUR AVERAGE</b>		Error includes scale factor of 2.8. See the ideogram below.			
2046 <sup>+17+67</sup> <sub>-16-15</sub>	1.8k	<sup>1</sup> ABLIKIM	20F	BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
1868 ± 8 <sup>+40</sup> <sub>-57</sub>	183k	ABLIKIM	19AQ	BES ±	$J/\psi \rightarrow K^+ K^- \pi^0$
2073 ± 94 <sup>+245</sup> <sub>-240</sub>	4289	<sup>2</sup> AAIJ	17C	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
1973 ± 8 ± 25		ASTON	87	LASS 0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
2020 ± 20		TIKHOMIROV 03	SPEC		40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1978 ± 40	241	BIRD	89	LASS -	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$

<sup>1</sup> Seen in  $\psi(2S)$  decay with branching ratio  $\psi(2S) \rightarrow K^\pm X \rightarrow K^+ K^- \eta = (7.0 \pm 0.5^{+3.7}_{-0.6}) \times 10^{-6}$ .

<sup>2</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of 5.4  $\sigma$ .



## $K_2^*(1980)$ WIDTH


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

**349<sup>+50</sup><sub>-30</sub> OUR AVERAGE** Error includes scale factor of 1.3. See the ideogram below.

408 <sup>+38</sup> <sub>-34</sub>	72 44	1.8k	<sup>1</sup> ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
272 $\pm$ 24 <sup>+50</sup> <sub>-15</sub>		183k	ABLIKIM	19AQBES $\pm$	$J/\psi \rightarrow K^+ K^- \pi^0$
678 $\pm$ 311 <sup>+1153</sup> <sub>-559</sub>		4289	<sup>2</sup> AAIJ	17C LHCB	$B^+ \rightarrow J/\psi \phi K^+$
373 $\pm$ 33 $\pm$ 60			ASTON	87 LASS 0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
180 $\pm$ 70			TIKHOMIROV 03	SPEC	40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
398 $\pm$ 47	241		BIRD	89 LASS -	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$

<sup>1</sup> Seen in  $\psi(2S)$  decay with branching ratio  $\psi(2S) \rightarrow K^\pm X \rightarrow K^+ K^- \eta = (7.0 \pm 0.5^{+3.7}_{-0.6}) \times 10^{-6}$ .

<sup>2</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of 5.4  $\sigma$ .



**$K_2^*(1980)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K^*(892)\pi$	possibly seen
$\Gamma_2$ $K\rho$	possibly seen
$\Gamma_3$ $K f_2(1270)$	possibly seen
$\Gamma_4$ $K\phi$	seen
$\Gamma_5$ $K\eta$	seen

 **$K_2^*(1980)$  BRANCHING RATIOS**

$\Gamma(K^*(892)\pi)/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>possibly seen</b>	GULER	11	BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$

$\Gamma(K\rho)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>possibly seen</b>	GULER	11	BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$

$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$   $\Gamma_2/\Gamma_1$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b><math>1.49 \pm 0.24 \pm 0.09</math></b>	ASTON	87	LASS	0 11 $K^- \rho \rightarrow \bar{K}^0 \pi^+ \pi^- n$

$\Gamma(K f_2(1270))/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>possibly seen</b>	TIKHOMIROV	03	SPEC $40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$

$\Gamma(K\phi)/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	4289	<sup>1</sup> AAIJ	17C	LHCB $B^+ \rightarrow J/\psi \phi K^+$

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $5.4 \sigma$ .

$\Gamma(K\eta)/\Gamma_{\text{total}}$   $\Gamma_5/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	1.8k	<sup>1</sup> ABLIKIM	20F	BES3 $\psi(2S) \rightarrow K^+ K^- \eta$
seen	116k	<sup>2</sup> CHEN	20A	BELL $D^0 \rightarrow K^- \pi^+ \eta$

<sup>1</sup> Seen decaying to  $K\eta$  in an amplitude analysis of  $\psi(2S) \rightarrow K^+ K^- \eta$ .

<sup>2</sup> From an amplitude analysis of the decay  $D^0 \rightarrow K^- \pi^+ \eta$  with a significance of  $17 \sigma$ .

 **$K_2^*(1980)$  REFERENCES**

ABLIKIM	20F	PR D101 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
CHEN	20A	PR D102 012002	Y.Q. Chen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
GULER	11	PR D83 032005	H. Guler <i>et al.</i>	(BELLE Collab.)
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>	
Translated from YAF 66 860.				

BIRD	89	SLAC-332	P.F. Bird	(SLAC)
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)

---